

Novel Mutations in Recessive Epidermolysis Bullosa Simplex

- receptors in cultured normal human epidermal keratinocytes. *Biochem Biophys Res Commun* 291:124–9
- Ito T, Ito N, Bettermann A, Tokura Y, Takigawa M, Paus R (2004) Collapse and restoration of MHC class-I dependent immune privilege: exploiting the human hair follicle as a model. *Am J Pathol* 164:623–34
- Maurer M, Fischer E, Handjiski B, Von Stebut E, Algermissen B, Bavandi A *et al.* (1997) Activated skin mast cells are involved in murine hair follicle regression (catagen). *Lab Invest* 77:319–32
- Müller-Röver S, Handjiski B, Van der Veen C, Eichmüller S, Foitzik K, McKay IA *et al.* (2001) A comprehensive guide for the accurate classification of murine hair follicles in distinct hair cycle stages. *J Invest Dermatol* 117:3–15
- Nakamura M, Sundberg JP, Paus R (2001) Mutant laboratory mice with abnormalities in hair follicle morphogenesis, cycling, and/or structure: annotated tables. *Exp Dermatol* 10:369–90
- Paus R, Hoffmann U, Eichmüller S, Czarnetzki BM (1994) Distribution and changing density of gamma-delta T cells in murine skin during the induced hair cycle. *Br J Dermatol* 130:281–9
- Paus R, van der Veen C, Eichmüller S, Kopp T, Hagen E, Müller-Röver S *et al.* (1998) Generation and cyclic remodeling of the hair follicle immune system in mice. *J Invest Dermatol* 111:7–18
- Paus R, Müller-Röver S, Van der Veen C, Maurer M, Eichmüller S, Ling G *et al.* (1999) A comprehensive guide for the recognition and classification of distinct stages of hair follicle morphogenesis. *J Invest Dermatol* 113:523–32
- Peters EMJ, Handjiski B, Kuhlmei A, Hagen E, Bielas H, Braun A *et al.* (2004) Neurogenic inflammation in stress-induced termination of murine hair growth is promoted by nerve growth factor. *Am J Pathol* 165:259–71
- Southall MD, Li T, Gharibova LS, Pei Y, Nicol GD, Travers JB (2003) Activation of epidermal vanilloid receptor-1 induces release of proinflammatory mediators in human keratinocytes. *J Pharmacol Exp Ther* 304:217–22
- Stenn KS, Paus R (2001) Controls of hair follicle cycling. *Phys Rev* 81:449–94
- Ständer S, Moormann C, Schumacher M, Buddenkotte J, Artuc M, Shpacovitch V *et al.* (2004) Expression of vanilloid receptor subtype 1 in cutaneous sensory fibers, mast cells, and epithelial cells of appendage structures. *Exp Dermatol* 13:129–39
- Szallasi A, Blumberg PM (1999) Vanilloid (Capsaicin) receptors and mechanisms. *Pharmacol Rev* 51:159–212
- Tominaga M, Caterina MJ, Malmberg AB, Rosen TA, Gilbert H, Skinner K *et al.* (1998) The cloned capsaicin receptor integrates multiple pain-producing stimuli. *Neuron* 21:531–43
- Veronesi B, Oortgiesen M, Carter JD, Devlin RB (1999) Particulate matter initiates inflammatory cytokine release by activation of capsaicin and acid receptors in a human bronchial epithelial cell line. *Toxicol Appl Pharmacol* 154:106–15

Novel Keratin 14 Mutations in Patients with Severe Recessive Epidermolysis Bullosa Simplex

Journal of Investigative Dermatology (2006) **126**, 1912–1914. doi:10.1038/sj.jid.5700312; published online 13 April 2006

TO THE EDITOR

Epidermolysis bullosa simplex (EBS) is the most common subtype, accounting for one-half of all epidermolysis bullosa cases (Pfundner *et al.*, 2005). It is clinically characterized by nonscarring blisters of the skin caused by little or no trauma, and morphologically by intra-epidermal blistering. The major subtypes of EBS result from mutations in either the keratin 5 (*KRT5*) or keratin 14 (*KRT14*) gene, whereas mutations in the gene for plectin (*PLEC1*) cause the rare forms, EBS with muscular dystrophy and EBS Ogna. The clinical spectrum of EBS ranges from mild blistering of the hands and feet (EBS Weber–Cockayne) to more generalized blistering (EBS Koebner, EBS Dowling–Meara, and EBS with mottled pigmentation). EBS, similarly to most of the keratin disorders identified in humans, is caused by domi-

nant missense mutations; however, patients with recessive EBS due to keratin mutations have been reported, representing about 5% of all EBS mutations (Porter and Lane, 2003).

In this study, we investigated two unrelated patients with severe neonatal blistering, both offspring of consanguineous, unaffected parents of Turkish (patient 1), respectively German (patient 2) origin. Indirect immunofluorescence (IIF) of the skin cryosections was performed as described (Hammami-Hausli *et al.*, 1998) with monoclonal antibodies anti-human keratin 5 (clone D5/16 B4, Dako, Hamburg, Germany) and keratin 14 (clone LL002, BioGenex, San Ramon, CA). Genomic DNA was extracted from peripheral blood samples collected from patients and their unaffected parents using the Qiagen Blood DNA Kit (Qiagen, Hilden, Ger-

many). Long-range polymerase chain reaction amplification of the *KRT14* gene was performed as described (Wood *et al.*, 2003) and direct sequencing in both directions was performed, using primers as published by Schuijnga-Hut *et al.* (2003) and an ABI prism 3100 automated sequencer (ABI, Darmstadt, Germany). The study was conducted according to the Declaration of Helsinki Principles, and the participants gave their written informed consent. The medical committee of the University of Freiburg approved all described studies.

Patient 1, a 2-year-old boy, showed blistering predominantly on hands and feet since birth (Figure 1a). In the course of the disease, bullae became rarer, occurred mechanically induced also on the head and trunk and healed without scarring. Patient 2, aged 1 year, showed at birth extensive blistering of the hands and feet (Figure 1b) and suffered from congenital pneumonia. Oral

Abbreviations: EBS, epidermolysis bullosa simplex; IIF, indirect Immunofluorescence; KRT, keratin




Figure 1. Phenotypes of the patients, indirect immunofluorescence staining of skin sections and mutations in the *KRT14* gene. (a) Blisters and erosions on the feet of patient 1 at the age of 2 years. (b) Patient 2 at 2 weeks of age showed severe blistering. (c) IIF with antibodies against keratin 5 showed intraepidermal splitting in the skin of patient 2. (d) IIF of control skin with antibodies against keratin 5. (e) IIF staining with antibodies against keratin 14 yielded a negative signal in patient's 1 skin. (f) IIF with antibodies against keratin 14 produced a positive signal in the basal keratinocyte layer in control skin. Bar = 100 μ m (given in (f) but refers to all images). (g) Partial sequence of *KRT14* exon 1, showing in the upper panel the mutation c.427delC (red arrow) and in the lower panel the wild-type sequence. (h) Partial sequence of *KRT14* exon 1, showing in the upper panel the mutation c.129dupC (red arrow) and in the lower panel the wild-type sequence.

blisters were present in the first days. Later, he developed blisters on arms, legs, and trunk after mechanical trauma. In both cases, IIF staining of skin sections revealed a split level through the basal cell layer (Figure 1c) and antibodies against keratin 14 yielded a negative staining (Figure 1e), suggesting keratin 14 defects in both cases. Mutation detection in *KRT14* revealed in patient 1 a homozygous one base pair deletion in exon 1, designated c.427delC (Figure 1g); this results in a frame shift starting with codon 143 and formation of a premature

termination codon, two codons downstream, p.L143fsX2. In patient 2, also in exon 1 of *KRT14*, a homozygous duplication, c.129dupC was disclosed (Figure 1h), leading to a frame shift starting with codon 44 and premature termination codon 38 codons downstream (p.S44fsX38). The parents were found to be heterozygous carriers of the respective mutations.

In both patients, the clinical picture was compatible with the diagnosis of EBS Koebner. Mucosal involvement and other symptoms are difficult to assess because of the very young age

and the short time they were available for observation. Absence of keratin 14 from the epidermis was caused by homozygous deletion/duplication mutations leading to premature termination codons. Important information was provided by IIF staining of a skin biopsy: first, intraepidermal blistering was evident and second, the negative signal with keratin 14 antibodies permitted the rapid identification of the candidate gene. Except for the two novel mutations described here, 11 different *KRT14* mutations associated with recessive EBS have been published before (Table 1): four with nonsense mutations, two with missense mutations, one with a splice site mutation, one compound heterozygous for nonsense and missense mutations, one with a deletion/insertion mutation, and two with deletion mutations. The less severely affected case reported by Batta *et al.* (2000) is the recessive case with the shortest expressible *KRT14* sequence to date, predicted to be truncated after only 30 amino acids (Porter and Lane, 2003; patient 6 in Table 1). In our patient 1, the truncation of keratin 14 is predicted to start after 143 amino acids, within the very beginning of the rod domain, and in patient 2 after 44 amino acids, within the head domain of keratin 14. Other authors described severely affected keratin 14 "knockout" patients (Table 1). However, no correlation between the position of the premature termination codon and the severity of the phenotype is obvious. Rather, these mutations are more likely to be associated with nonsense-mediated messenger RNA decay. Chan *et al.* (1994) suggested that complete absence of a keratin is less detrimental than the disrupted filament assembly and aggregate formation caused by dominant-negative missense mutations. The present findings and other reports in the literature agree with this hypothesis in that homozygous missense mutations lead to milder phenotypes with blistering only in the extremities (patients 1, 10, and 11 in Table 1), whereas nonsense mutations are associated with more generalized blistering (patients 2, 3, 4, 5, 7, 8, and 9 in Table 1).

Table 1. KRT14 mutations in recessive EBS

No.	Phenotype, age	Mutations	References
1.	Blistering affecting the lateral and dorsal aspects of the feet and plants, two patients aged 20 and 12 y	p.E144A/p.E144A	Hovnanian <i>et al.</i> (1993)
2.	Severe generalized blistering, 5 y	c.313_314delGC/ c.313_314delGC	Rugg <i>et al.</i> (1994)
3.	Generalized blistering, 29 m	p.Y204X/p.Y204X	Chan <i>et al.</i> (1994)
4.	Severe generalized blistering, mucous membranes occasionally affected; family with 4 patients aged 74, 67, 47, 34 y	c.1842-2A>C/c.1842-2A>C	Jonkman <i>et al.</i> (1996)
5.	Generalized blistering, occasional oral blisters, siblings of 7 and 6 y	p.W305X/p.W305X	Corden <i>et al.</i> (1998)
6.	Mild generalized blistering, 18 m	c.92delT/c.92delT p.I31fs86X	Batta <i>et al.</i> (2000)
7.	Generalized skin blistering and mild nail involvement and involvement of mucous membranes, 8 y	c.744delC/insAG/ c.744delC/ insAG p.Y248X	Landschuetzer <i>et al.</i> (2003)
8.	Widespread blistering and oral mucosal involvement, anaemia and failure to thrive in the first year of life, NA	p.Q396X/p.Q396X	Ciubotaru <i>et al.</i> (2003)
9.	Widespread blistering over palms and soles and oral and genital mucosa, NA	p.W305X/p.W305X	Ciubotaru <i>et al.</i> (2003)
10.	Blisters over palms and soles, NA	p.Q396X/p.R388H	Ciubotaru <i>et al.</i> (2003)
11.	Blisters over hands and feet, 4 y	p.R134C/p.R134C	Indelman <i>et al.</i> (2005)
12.	Blisters on hands and feet, later generalized, 2 y	c.427delC/c.427delC p.L143fsX2	This study
13.	Severe blistering of hands and feet, oral erosions, widespread induced blisters, 1 y	c.129dupC/c.129dupC p.S44fsX38	This study

m, months; NA, not available; y, years.

CONFLICT OF INTEREST

The authors state no conflict of interest.

ACKNOWLEDGMENTS

We thank the patients and their families and the partners of the German *Netzwerk Epidermolysis bullosa* (www.netzwerk-eb.de) for referring patients for mutation analysis. This work was supported in part by the EB-Network grant from the German Ministry for Research and Education (BMBF).

**Cristina Has¹, Yow-Ren Chang¹,
Andreas Volz¹, Doris Hoeping¹,
Jürgen Kohlhas² and Leena
Bruckner-Tuderman¹**

¹Department of Dermatology, University of Freiburg, Freiburg, Germany and ²Institute of Human Genetics and Anthropology, University of Freiburg, Freiburg, Germany.
E-mail: crishas@haut.ukl.uni-freiburg.de

REFERENCES

- Batta K, Rugg EL, Wilson NJ, West N, Goodyear H, Lane EB *et al.* (2000) A keratin 14 'knockout' mutation in recessive epidermolysis bullosa simplex resulting in less severe disease. *Br J Dermatol* 143:621-7
- Chan Y, Anton-Lamprecht I, Yu QC, Jackel A, Zabel B, Ernst JP *et al.* (1994) A human keratin 14 "knockout": the absence of K14 leads to severe epidermolysis bullosa simplex and a function for an intermediate filament protein. *Genes Dev* 8:2574-2587
- Ciubotaru D, Bergman R, Baty D, Indelman M, Pfendner E, Petronius D *et al.* (2003) Epidermolysis bullosa simplex in Israel. *Arch Dermatol* 139:498-505
- Corden L, Mellerio J, Gratian MJ, Eady AJ, Harper JL, Lacour M *et al.* (1998) Homozygous nonsense mutation in helix 2 of K14 causes severe recessive epidermolysis bullosa simplex. *Hum Mutat* 11:279-85
- Hammami-Hauasli N, Schumann H, Raghunath M, Kilgus O, Luthi U, Luger T *et al.* (1998) Some, but not all, glycine substitution mutations in COL7A1 result in intracellular accumulation of collagen VII, loss of anchoring fibrils, and skin blistering. *J Biol Chem* 273:19228-34
- Hovnanian A, Pollack E, Hilal L, Rochat A, Prost C, Barrandon Y *et al.* (1993) A missense mutation in the rod domain of keratin 14 associated with recessive epidermolysis bullosa simplex. *Nat Genet* 3:327-32
- Indelman M, Bergman R, Sprecher E (2005) A novel recessive missense mutation in KRT14 reveals striking phenotypic heterogeneity in epidermolysis bullosa simplex. *J Invest Dermatol* 124:272-4
- Jonkman MF, Heeres K, Pas HH, Van Luyn MJA, Elema JD, Corden LD *et al.* (1996) Effects of keratin 14 ablation on the clinical and cellular phenotype in a kindred with recessive epidermolysis bullosa simplex. *J Invest Dermatol* 107:764-9
- Landschuetzer CM, Klausegger A, Pohla-Gubo G, Hametner R, Richard G, Uitto J *et al.* (2003) A novel homozygous nonsense deletion/insertion mutation in the keratin 14 gene (Y248X; 744delC/insAG) causes recessive epidermolysis bullosa simplex Köbner. *Clin Exp Dermatol* 28:77-9
- Pfendner EG, Sadowski SG, Uitto J (2005) Epidermolysis bullosa simplex: recurrent and *de novo* mutations in the KRT5 and KRT14 genes, phenotype/genotype correlations and implications for genetic counselling and prenatal diagnosis. *J Invest Dermatol* 125:239-43
- Porter R, Lane B (2003) Phenotypes, genotypes and their contribution to understanding keratin function. *Trends Genet* 19:278-85
- Rugg EL, McLean WH, Lane EB, Pitera R, McMillan JR, Dopping-Hepenstal PJ *et al.* (1994) A functional "knockout" of human keratin 14. *Genes Dev* 8:2563-73
- Schuilenga-Hut PH, Vlies P, Jonkman MF, Waanders E, Buys CH, Scheffer H (2003) Mutation analysis of the entire keratin 5 and 14 genes in patients with epidermolysis bullosa simplex and identification of novel mutations. *Hum Mutat* 21:447
- Wood P, Baty DU, Lane EB, McLean WH (2003) Long-range polymerase chain reaction for specific full-length amplification of the human keratin 14 gene and novel keratin 14 mutations in epidermolysis bullosa simplex patients. *J Invest Dermatol* 120:495-7